



## Diflufenican

Diflufenican is a selective contact and residual herbicide, absorbed principally by the shoots of germinating seedlings, with limited translocation. Uses Applied at 125-250 g/ha pre- or early post-emergence in autumn-sown wheat and barley, to control grass and broad-leaved weeds, particularly *Galium*, *Veronica* and *Viola* spp. Normally used in combination with isoproturon or other cereal herbicides.



**Action:** \* Selective contact

\* Pre- or early post-emergence

**CAS No.:** 83164-33-4

**Technical Grade:** 98% min, 99% min

**Formulations:** 50%SC, 50%WP

**Packaging<sup>1</sup>:** \* 25kg drum (TC)

\* 1kg bag (WP)

\* 1L bottle (SC)

<b>Crops</b>	<b>Target Weeds</b>	<b>Dose (a.i. g/ha)</b>
<i>Wheat</i>	Broad-leaved weeds and gramineae weeds	125~250

<sup>1</sup>NOTE: Upon request we provide customized packing

This document was created by Unregistered Version of Word to PDF ConverterCopyright Owner©

2010 **FlagChem**

- Changfenghe Road, Nanjing Chemical Industrial Park, Luhe, Nanjing, China 210047
- Tel: 86-25-58375015 Fax: 86-25-58375450
- E-mail: [sales@flagchem.com](mailto:sales@flagchem.com) Website: [www.flagchem.com](http://www.flagchem.com)