



## Cyprodinil

Cyprodinil is a foliar fungicide use in cereals, grapes, pome fruit, stone fruit, strawberries, vegetables, field crops and ornamentals; and as a seed dressing on barley. It is used to inhibit the biosynthesis of methionine and the secretion of fungal hydrolytic enzymes. There is no cross resistance to benzimidazoles, carbamates, dicarboximides, imidazoles, morpholines, quinolines, strobilurins or triazoles.



**Action:** \*Systemic product

\* Foliar application

**CAS No.:** 121552-61-2

**Technical Grade:** 95% min.

**Packaging<sup>1</sup>:** \* 25kg drum

Crops	Target Diseases	Dose (a.i. g/ha)
Cereals	<i>Tapesia yallundae</i>	150-750 g/ha

<sup>1</sup>NOTE: Upon request we provide customized packing

This document was created by Unregistered Version of Word to PDF ConverterCopyright Owner©  
2010 **FlagChem**

- Changfenghe Road, Nanjing Chemical Industrial Park, Luhe, Nanjing, China 210047
- Tel: 86-25-58375015 Fax: 86-25-58375450
- E-mail: [sales@flagchem.com](mailto:sales@flagchem.com) Website: [www.flagchem.com](http://www.flagchem.com)